Lake Erie Bathymetry and Physiographic Features



LAKE ERIE WESTERN BASIN ISLANDS

1 - Turtle Island	9 - Chick Island
2 - West Sister Island	10 - Sugar Island
3 - Middle Sister Island	11 - Rattlesnake Island
4 - North Harbour Island	12 - Green Island
5 - East Sister Island	13 - Mouse Island
6 - Hen Island	14 - Ballast Island
7 - Little Chicken Island	15 - Middle Island
8 - Big Chicken Island	16 - Johnson's Island

LAKE ERIE CANADIAN "POINTS"

1 - Patrick Point	11 - Mohawk Point
2 - Plum Point	12 - Grabell Point
3 - Turkey Point	13 - Morgans Point
4 - Peacock Point	14 - Rathfon Point
5 - Hoover Point	15 - Sugarloaf Point
6 - Featherstone Point	16 - Cassady Point
7 - Evans Point	17 - Pine Crest Point
8 - Blott Point	18 - Shisler Point
9 - Low Point	19 - Sunrise Point
10 - Rock Point	20 - Windmill Point

0 10 20 30 40

Miles

Bathymetry Legend

Minimum Depth — Shallower — ≈0-3 ft (0-1 m)

Average Depth — 62 ft (19 m)

Maximum Depth — Deeper — 210 ft (64 m)

GIS Data Sources

Lake Erie Bathymetry - National Oceanic and Atmospheric Administration (NOAA), 2004
Lake Erie and Lake St. Clair Shoreline - NOAA, 2004

This map is a product of the Ohio Department of Natural Resources (ODNR) Office of Coastal Management and is intended to provide general information only.

coastal.ohiodnr.gov Map created: 11/14